

Progress on indium and barium single ion optical frequency standards

Jeff A. Sherman, William Trimble, Steven Metz, Warren Nagourney, and Norval Fortson

Abstract—We report progress on $^{115}\text{In}^+$ and $^{137}\text{Ba}^+$ single ion optical frequency standards using all solid-state sources. Both are free from quadrupole field shifts and together enable a search for drift in fundamental constants.

I. INTRODUCTION

IN single ion optical frequency standards the narrow transitions of weakly allowed lines serve as high quality references largely free from shifts and perturbations such as the first-order Doppler effect. Other research groups have demonstrated precise control or near-elimination of most effects, such as the second-order Doppler shift, quadratic Stark shifts, Zeeman shifts, and ac Stark shifts from black-body radiation for a diverse set of ion species (In^+ [1], Hg^+ [2], Yb^+ [3], [4], Sr^+ [5], Ca^+ [6], and proposed in Tl^+ [7]). However, one important effect that remains to be suitably controlled at the Hz level is the quadrupole Stark shift—that is, the shift arising from coupling of the trapped ion's atomic quadrupole moment to stray dc electric field gradients unavoidably present in the trap region.

Here we propose two single ion optical frequency standards that are inherently free from this effect and are under active development. Solid-state lasers for all the required transitions are commercially available. In addition to being viable candidates for absolute frequency standards, their construction can also yield a search for drift in fundamental physical constants.

II. INDIUM ION FREQUENCY STANDARD

We trap single indium ions in a twisted wire Paul-Straubel trap with 700 volts of 10 MHz rf. We then cool the ions to the Lamb-Dicke regime using 230.6 nm light resonant with the $^1S_0 \leftrightarrow ^3P_1$ ($F = 11/2$) transition. This radiation is generated by frequency quadrupling a 922 nm external cavity diode laser/tapered amplifier system in resonant build-up cavities containing KNbO_3 and BBO nonlinear crystals. The 3P_0 excited state is long-lived ($\tau = 195(8)$ ms [1]) giving the $^1S_0 \leftrightarrow ^3P_0$ transition a natural linewidth of ~ 1 Hz and a quality factor of $Q = 1.5(10)^{15}$. We create 236.5 nm clock light by frequency quadrupling the output of a monolithic Nd:YAG laser operating at 946 nm. This laser, manufactured by InnoLight GmbH, is a non-planar ring oscillator with a nominal free-running linewidth of 2 kHz. Similar laser systems have been narrowed below 1 Hz [8] by Pound-Drever-Hall locking to an isolated high finesse reference cavity.

Excitation of the clock transition is detected via the absence of cooling fluorescence when the ion is ‘shelved’ in




Fig. 1

RELEVANT ATOMIC LEVELS AND TRANSITIONS FOR $^{115}\text{In}^+$ AND $^{137}\text{Ba}^+$ FREQUENCY STANDARDS. ALL THE WAVELENGTHS ARE ACCESSIBLE WITH SOLID STATE LASERS AS DISCUSSED IN THE TEXT.

the metastable 3P_0 state. A software servo system detects and corrects for long-term drifts in the clock laser while it is stabilized over short times using a stable reference cavity or other absolute reference, such as a microwave standard and femtosecond frequency comb.

An important feature of $^{115}\text{In}^+$ is its immunity to quadrupole Stark shifts due to both ground and excited clock states having zero electronic angular momentum $J = J' = 0$. From Ref. [9], quadrupole Stark shifts take the form

$$H_Q = \nabla \mathbf{E}^{(2)} \cdot \boldsymbol{\Theta}^{(2)} \quad (1)$$

where the tensor $\nabla \mathbf{E}^{(2)}$ describes the electric field gradient at the site of the ion and $\boldsymbol{\Theta}^{(2)}$ are the set of irreducible second-rank tensors. In the IJ -coupling approximation, shifts of the clock frequency scale with the reduced matrix elements

$$\langle n(IJ)F || \boldsymbol{\Theta}^{(2)} || n(IJ)F \rangle \propto \begin{Bmatrix} J & 2 & J \\ F & I & F \end{Bmatrix} \langle nJJ | \boldsymbol{\Theta}_0^{(2)} | nJJ \rangle \quad (2)$$

where the $6j$ -symbol and the atomic quadrupole moment vanish for $J = 0$.

	$^{115}\text{In}^+$	$^{137}\text{Ba}^+$
Clock frequency (cm^{-1})	42275.995	4873.720
Clock state lifetime (s)	0.2 [1]	80 [15]
Quality factor $\nu/\Delta\nu$	$1.5(10)^{15}$	$7(10)^{16}$
Zeeman shift	224 Hz/G [1]	-11 kHz/G ²
Clock drift (Hz/yr) (if $\dot{\alpha}/\alpha = 10^{-15} \text{ yr}^{-1}$)	0.23 [17]	0.35 [18]

TABLE I
SELECTED PROPERTIES OF THE TWO SINGLE ION FREQUENCY
STANDARDS DISCUSSED HERE.

III. BARIUM ION FREQUENCY STANDARD

Our barium ion trap [10] operates at a similar frequency and strength as the indium trap but is constructed entirely from non-magnetic tungsten and tantalum. Passive shielding has achieved $\leq 10 \mu\text{G}$ magnetic field stability measured using the ion [11]. We cool on the $493 \text{ nm } 6S_{1/2} \leftrightarrow 6P_{3/2}$ transition using a frequency doubled 986 nm diode laser/tapered amplifier system while also applying a 650 nm diode laser which empties the long-lived $5D_{3/2}$ state. Currently, a static magnetic field ($B \sim 1.6 \text{ G}$) aligned along both lasers eliminates dark states [12] but for clock applications a low magnetic field is desirable; polarization modulation will accomplish dark state quenching [13]. High intensity filtered light-emitting diodes perform shelving to and deshelling from the $5D_{5/2}$ state and could ultimately be replaced by a $1.76 \mu\text{m}$ diode laser acting on the $6S_{1/2} \leftrightarrow 5D_{5/2}$ electric-quadrupole transition previously investigated as a clock candidate [14].

The $\text{Ba}^+ 5D_{3/2}$ state lifetime ($\tau \sim 80 \text{ s}$ [15], [16]) is among the longest-lived of successfully cooled ions, giving the $6S_{1/2} \leftrightarrow 5D_{3/2}$ clock transition at 2051 nm a quality factor $Q = \nu/\Delta\nu > 10^{16}$. A diode pumped Tm,Ho:YLF solid state laser manufactured by CLR Photonics produces $> 80 \text{ mW}$ at 2051 nm with a free-running linewidth we have determined to be $\sim 10 \text{ kHz}$. We have detected the transition in $^{138}\text{Ba}^+$.

$^{137}\text{Ba}^+$ shares the advantage with In^+ of having vanishing shifts due to quadrupole Stark shifts. The atomic quadrupole moment in Eq. 2 vanishes for the ground state $6S_{1/2}(F=2)$ and the $6j$ -symbol in Eq. 2 vanishes for the excited clock state $5D_{3/2}(F=0)$. Small shifts likely remain due to configuration mixing and coupling to the nuclear moment but these perturbations are expected to be much less than 1 mHz.

IV. A SEARCH FOR DRIFT IN FUNDAMENTAL CONSTANTS

Besides applications in navigation, long baseline interferometry, and detection of drift in pulsar periodicity, optical atomic frequency standards enable laboratory searches for drift in fundamental constants [19]. Cosmological observations [20] as well the analysis of the Oklo natural nuclear reactor [21] suggest drift in the fine-structure constant. Since relativistic corrections to clock transition frequencies of different atomic species depend differently on the constant α ,

a linear drift of α over time will be detected as an accumulated error between two or more atomic optical frequency standards. Table I shows the estimated clock frequency change that would occur in one year assuming a fractional change of $\dot{\alpha}/\alpha = 10^{-15} \text{ yr}^{-1}$ using latest theoretical results [17], [18].

REFERENCES

- [1] T. Becker, *et al.*, “High-resolution spectroscopy of a single In^+ ion: Progress towards an optical frequency standard,” *Phys. Rev. A*, vol. 63, no. 5, p. 051802, May 2001.
- [2] S. A. Diddams, *et al.*, “An Optical Clock Based on a Single Trapped $^{199}\text{Hg}^+$ Ion,” *Science*, vol. 293, pp. 825–828, Aug. 2001.
- [3] P. J. Blythe, *et al.*, “Subkilohertz absolute-frequency measurement of the 467-nm electric octupole transition in $^{171}\text{Yb}^+$,” *Phys. Rev. A*, vol. 67, no. 2, p. 020501, Feb. 2003.
- [4] C. Tamm, D. Engelke, and V. Bühner, “Spectroscopy of the electric-quadrupole transition $^2S_{1/2}(F=0) \rightarrow ^2D_{3/2}(F=2)$ in trapped $^{171}\text{Yb}^+$,” *Phys. Rev. A*, vol. 61, no. 5, p. 053405, May 2000.
- [5] G. P. Barwood, *et al.*, “Subkilohertz comparison of the single-ion optical-clock $^2S_{1/2} \rightarrow ^2D_{5/2}$ transition in two $^{88}\text{Sr}^+$ traps,” *Phys. Rev. A*, vol. 59, p. 3178, May 1999.
- [6] C. Champenois, *et al.*, “Evaluation of the ultimate performances of a Ca^+ single-ion frequency standard,” *Phys. Lett. A*, vol. 331, pp. 298–311, Oct. 2004.
- [7] J. R. Torgerson, “Method for precision test of fine structure constant variation with optical frequency references,” Dec. 2000, e-print: physics/0012054.
- [8] S. A. Webster, M. Oxborrow, and P. Gill, “Subhertz-linewidth Nd:YAG laser,” *Opt. Lett.*, vol. 29, no. 13, pp. 1497–99, jul 2004.
- [9] W. M. Itano, “External-field shifts of the $^{199}\text{Hg}^+$ optical frequency standard,” *J. Research NIST*, vol. 105, no. 6, pp. 829–37, 2000, e-print: physics/0010056v2.
- [10] T. W. Koerber, *et al.*, “Radio frequency spectroscopy with a trapped Ba^+ ion: recent progress and prospects for measuring parity violation,” *J. Phys. B*, vol. 36, no. 3, pp. 637–648, 2003.
- [11] J. Sherman, *et al.*, “Precision measurement of light shifts in a single trapped Ba^+ ion,” 2005, submitted to *Phys. Rev. Lett.*, e-print: physics/0502113.
- [12] D. J. Berkeland and M. G. Boshier, “Destabilization of dark states and optical spectroscopy in Zeeman-degenerate atomic systems,” *Phys. Rev. A*, vol. 65, no. 3, p. 033413, 2002.
- [13] R. G. DeVoe and C. Kurtsiefer, “Experimental study of anomalous heating and trap instabilities in a microscopic ^{137}Ba ion trap,” *Phys. Rev. A*, vol. 65, no. 6, p. 063407, 2002.
- [14] N. Yu, *et al.*, “Stark shift of a single barium ion and potential application to zero-point confinement in a rf trap,” *Phys. Rev. A*, vol. 50, no. 3, pp. 2738–41, 1994.
- [15] N. Yu, W. Nagourney, and H. Dehmelt, “Radiative Lifetime Measurement of the Ba^+ metastable $D_{3/2}$ State,” *Phys. Rev. Lett.*, vol. 78, pp. 4898–4901, June 1997.
- [16] A. A. Madej and J. D. Sankey, “Quantum jumps and the single trapped barium ion: Determination of collisional quenching rates for the $5d^2D_{5/2}$ level,” *Phys. Rev. A*, vol. 41, no. 5, pp. 2621–30, 1990.
- [17] E. J. Angstmann, V. A. Dzuba, and V. V. Flambaum, “Relativistic effects in two valence electron atoms and ions and the search for variation of the fine structure constant,” *Phys. Rev. A*, vol. 70, no. 1, p. 014102, July 2004.
- [18] V. A. Dzuba and V. V. Flambaum, “Atomic optical clocks and search for variation of the fine-structure constant,” *Phys. Rev. A*, vol. 61, no. 3, p. 034502, Mar. 2000.
- [19] S. Bize, *et al.*, “Testing the stability of fundamental constants with the $^{199}\text{Hg}^+$ single-ion optical clock,” *Phys. Rev. Lett.*, vol. 90, no. 15, p. 150802, Apr. 2003.
- [20] M. T. Murphy, J. K. Webb, and V. V. Flambaum, “Further evidence for a variable fine-structure constant from Keck/HIRES QSO absorption spectra,” *Mon. Not. R. Astron. Soc.*, vol. 345, no. 2, pp. 609–38, Oct. 2003.
- [21] S. K. Lamoreaux and J. R. Torgerson, “Neutron moderation in the Oklo natural reactor and the time variation of alpha,” *Phys. Rev. D*, vol. 69, no. 12, p. 121701, 2004.